

Cosmetic and/or dermatological use of a composition containing at least one oxidation-sensitive hydrophilic active stabilised by at least one polymer or copolymer of maleic anhydride

Publication number: EP1374849

Publication date: 2004-01-02

Inventor: BIATRY BRUNO (FR)

Applicant: OREAL (FR)

Classification:






- International: A61K8/00; A61K8/02; A61K8/06; A61K8/11; A61K8/30; A61K8/31; A61K8/33; A61K8/34; A61K8/35; A61K8/36; A61K8/365; A61K8/368; A61K8/37; A61K8/40; A61K8/41; A61K8/42; A61K8/44; A61K8/46; A61K8/49; A61K8/55; A61K8/58; A61K8/60; A61K8/63; A61K8/67; A61K8/68; A61K8/72; A61K8/73; A61K8/81; A61K8/92; A61K8/96; A61K8/97; A61K8/98; A61K31/375; A61K45/00; A61K47/32; A61P17/00; A61P17/16; A61P39/06; A61P43/00; A61Q5/00; A61Q5/08; A61Q19/00; A61Q19/02; A61Q19/08; C07D307/62; A61K47/32; A61K8/00; A61K8/02; A61K8/04; A61K8/11; A61K8/30; A61K8/72; A61K8/92; A61K8/96; A61K31/375; A61K45/00; A61P17/00; A61P39/00; A61P43/00; A61Q5/00; A61Q5/08; A61Q19/00; A61Q19/02; A61Q19/08; C07D307/00; (IPC1-7): A61K7/48; A61K7/00; A61K7/06

- European: A61K8/11C; A61K8/34F; A61K8/49H; A61K8/67H; A61K8/81M; A61K8/81R4; A61Q5/08; A61Q19/00; A61Q19/02; A61Q19/08

Application number: EP20030291008 20030424






Priority number(s): FR20020007638 20020620

Also published as:

 EP1374853 (A1)
 EP1374852 (A1)
 EP1374851 (A1)
 EP1374850 (A1)
 JP2004067677 (A)

more >>

Cited documents:

 US6103267
 FR2816316
 EP0815847
 EP0380367
 EP1151741

more >>

[Report a data error here](#)

Abstract of EP1374849

In a cosmetic composition containing an aqueous phase, a combination of at least one oxidation sensitive hydrophilic active principle (a) and at least one maleic anhydride copolymer (b) are used as depigmenting agents. (a) Is ascorbic acid or its derivatives. (b) Comprises at least one maleic anhydride comonomers and at least one of e.g. vinyl acetate, vinyl alcohol or styrene. In a cosmetic composition containing an aqueous phase, a combination of at least one oxidation sensitive hydrophilic active principle (a) and at least one maleic anhydride copolymer (b) are used as depigmenting agents. (a) Is ascorbic acid or its derivatives. (b) Comprises at least one maleic anhydride comonomers and at least one of vinyl acetate, vinyl alcohol, vinyl pyrrolidone, olefins containing 2-20C atoms or styrene.

Data supplied from the esp@cenet database - Worldwide